ACL Reconstruction Rehabilitation Program

The Gundersen Health System Sports Medicine ACL Reconstruction Rehabilitation Program is an evidence-based and soft tissue healing dependent program allowing patients to progress to vocational and sports-related activities as quickly and safely as possible. Individual variations will occur depending on surgical technique and the patient's response to treatment.

Early utilization of NMES highly recommended and encouraged with 2x4", ideally 3x5" stim pads.

Early isolated OKC resisted knee extension should be implemented depending on patient tolerance- ie no increase in PF pain/joint effusion.

If a meniscus repair is performed in conjunction with the ACL reconstruction, follow meniscus repair WB restrictions (stable or unstable), ROM 0-90 for 2 wk, no squatting >90 for 4 months, as can be seen in meniscus repair protocol.

If a hamstring/gracilis autograft is utilized, avoid isolated hamstring strengthening for 6 weeks. If a patellar tendon graft is utilized, work on patella mobilizations to prevent excessive scaring. 21771 If an allograft is utilized, patients may need to be cautioned not to advance too quickly as post-operative pain may be less.

Please contact us at 1-800-362-9567 ext. 58600 if you have questions or concerns.

Phase I: 0-6 weeks	Immediate post op maximum protection phase
Goals	 Protect surgical graft Minimize knee joint effusion Restore full motion ASAP, emphasizing extension Encourage quadriceps function Prevent negative effects of immobilization Normalization of walking with good heel-toe pattern Not all patients will utilize a post-operative brace.
Biace	 • Not all patients will utilize a post-operative brace. • wks 0-1: Locked for ambulation and sleeping • wks 1+: set per available ROM, full if able, unlocked for ambulation when good quadriceps control and ext ROM full • wk 4: D/C brace
ROM	 Full extension and full flexion immediately post op without restriction, emphasis on extension initially with gradual progression of flexion Goal is full ROM by 6 wk
WB	 wk 0-1: WBAT with brace locked into extension wk 1-4: WBAT with brace unlocked if good quadriceps control and knee extension ROM. Discontinue crutches when can ambulate with normal heel-to-toe pattern.
Precautions	 If hamstring/gracilis autograft, no isolated resistance to knee flexion initially. Start graded isometrics at wk 4 within tolerance. Progress to isotonics at wk 6. Also apply ice to posterior knee to minimize muscle spasm. Encourage AROM and WB to promote healing, prevent atrophy of soft tissue and bone, prevent a decrease in collagen content, and to align fibroblast and collagen fibrils. Emphasis on regaining extension ROM ASAP to prevent arthrofibrosis and decrease stress to the PF joint during ambulation. Avoid descending stair reciprocally until adequate quadriceps control and lower extremity alignment Avoid twisting and pivoting motions for 6-8 weeks to minimize shear forces to the healing graft.



Modalities • Cryotherapy 15 minutes in duration 3x/day • IFC for pain/effusion if needed • NMES quadriceps ASAP in varying positions using 2x4" but ideally 3x5" pads Long sitting QS/SLR/SAQ Short sitting LAQ isometrics into strap vs isotonics with resistance Standing TKE with TB or CC resistance Updated 11/2023 **Treatment** Active warm-up (Bike AAROM progress to Bike with resistance, Nu Step) Recommendations • Stretching to attain full extension with gradual progression of flexion. Goal of full ROM BEFORE wk 6. Emphasis on full return of knee extension ASAP. Low-load long duration stretching for extension with heat if needed (1st TERT= Total End Range Time) Manual stretching for extension with overpressure / recurvatum Patellar mobilizations Guidelines for PROM / AAROM / AROM progression based on Manual stretching into flexion (initially limited by knee joint effusion) tolerance wk 4: WB stretch on leg press for knee flexion ROM • Flexibility exercises for hamstring, gastoc-soleus • Scar tissue massage • Consider personalized blood flow restriction therapy if available. • Therapeutic exercises. Gentle strengthening protecting the surgical graft. Encourage quadriceps activation and strength. Early isolated OKC resisted knee extension can be implemented depending on patient tolerance- ie no increase in PF pain/joint effusion. Exercise in a pain-free manner. Avoid dynamic valgus during strengthening and functional activities (focus on hip abductor and external rotator strengthening). Incorporate total leg strengthening and balance / proprioception exercises. Work on gait drills (step-overs, march walk). CKC TKEs Hip 4 way SLR, sidelying hip ER Gastroc soleus strengthening Hamstring OKC isotonics 0-90 deg in seated position **Quadriceps OKC knee extension per patient tolerance** CKC exercises: Heel raises, weight shifts, leg press and wall squats (0-60 deg) wk 2: Leg press, wall squats (0-90 deg), lateral step-overs step-ups, BW squats, retro TM walking for knee extension forward TM walking for gait training wk 3: Partial lunges front and lateral, leg press 2:1, loaded squats as tolerated wk 4: Elliptical Runner, leg press 2:1 and 1:1, deadlifts wk 5: Resisted sidestep with T-band, Bosu partial squats 0-60 deg Total leg strengthening Balance / Proprioception training: Double leg progress to single leg, static progressing to dynamic activities. Perturbation exercises CV conditioning / Core Stability • IFC for pain/effusion, NMES for quadriceps activation and control as needed Ice (in stretch for extension if needed) 2nd TERT HEP for 3rd TERT

Phase II: 6-12 weeks	Moderate protective phase
Goals	Minimize knee joint effusion
	Full ROM by 6-8 weeks
	Gradual progression of exercises for strengthening, stretching, and balance
	Implement low level foot placement drills working on control
	Ability to squat with symmetry to 90 degrees and single leg squat with good
	knee flexion, control of any dynamic valgus, and without pelvic drop
ROM /	Full by 6 wk.
Brace	Knee sleeve may be utilized depending on patient activities
Modalities	Cryotherapy 15 minutes in duration 1-2x/day
	IFC for pain/effusion if needed. NMES quadriceps if needed
Precautions	Avoid overloading the fixation site by utilizing low amplitude low velocity
	movements.
	 Avoid quick twisting and pivoting motions for 10-12 wks to minimize shear
	forces.
	Implement low level foot placement focus on control at week 9.
Treatment	Active warm-up: Bike with resistance, Nu Step, Treadmill walking
Recommendations	Stretching for full extension and flexion as needed.
	Low-load long duration stretching with heat if needed
	(1st TERT= Total End Range Time)
Guidelines for	Manual stretching for extension and/or flexion
progression	Leg press stretch for flexion
based on tolerance	Flexibility exercises as needed
	Therapeutic exercises: Focus on N-M control and strengthening
	exercises. Avoid dynamic valgus during strengthening and functional
	activities. Incorporate total leg strengthening, focus on hip/glutes,
	quadriceps, and hamstring. Progress with balance / proprioception exercises.
	Correct asymmetrical loading patterns
	Total leg strengthening and CV conditioning Hip and core strengthening to prevent knee valgus
	Hamstrings isotonics prone 0-90 deg.
	Quadriceps isotonics OCK knee extension
	Balance / Proprioception training: Single leg stance activities
	static progressing to dynamic activities. Perturbation
	exercises
	CKC exercises: Leg press 1:1, step-ups/step downs,
	squats progression double to single leg, split squats
	lunge progression, deadlifts, sidestep/sideshuffle with SPRI
	wk 8: Hamstring curls with physio ball
	Balance exercises: add in external focus of attention (ball
	catch, plyo back throws)
	Isokinetic quadriceps/hamstrings
	VSRP 180-300 deg/sec sub-max to max
	Wk 10: Isokinetic quadriceps/hamstring
	VSRP 60-300 deg/sec sub-max to max
	 Low level foot placement drills starting at wk 9 IFC for pain/effusion / NMES for quadriceps activation and control as needed
	 Ice (in stretch if needed) 2nd TERT
	HEP for 3 rd TERT if needed
Indonandant	
Independent strengthening	wk 12: Can progress to independent strengthening program with monthly or bi-monthly visits if good ROM, minimal effusion, full MMT, able to SL squat,
on onguiening	pass y-balance.
	pass y-balance.
	I



Phase III: 12-24 wks	Advanced Strengthening and Functional Activities
(3-6 months)	
Make sure patient is enrolled in MyCare for IKDC survey (6M, 9M, 1Y, 2 Y, 5Y)	 Progress muscle strength, endurance, and balance activities. Ideally 3x/week of exercises at a fitness center, step-down, or home program. Progress to higher level activities depending on functional demands and MD approval Address fear avoidance beliefs by graded exercise progression, cuing, positive reinforcement, referral if necessary. Initiate a return to running program at 3-4 months if passes criteria and has no compensations with running pattern. Initiate working on landing mechanics and control at 4-5 months if passes criteria on the following page Progress agility drills at 4-5 months
Brace	Your MD may recommend a knee sleeve or functional brace to be used until 12 months from your surgery for higher level activities
Modalities	Cryotherapy 15 minutes 1x/day or after strenuous activity
Treatment	Active warm-up:
Recommendations	Continue with stretching and flexibility exercises as needed
	Strengthening and endurance exercises: Focus on strengthening and N-M
Poturn to Dunning	control activities. Advance as tolerated with emphasis on functional
Return to Running Benchmarks:	strengthening. Focus on soft landing with knee flexion, no medial
4 months	collapse/knee valgus, and postural control. Progress with balance /
Passes testing criteria -	proprioception exercises. Progress to working on landing mechanics and some agility drills as appropriate. Correct assymetrical loading patterns
See next page	Total leg strengthening: hip/quadriceps/hamstring
	Hip strengthening – neuromuscular control to prevent knee valgus
	Core strengthening – prevent frontal plane trunk lean during landing/SLS
	Hamstring full ROM isotonics
Return to Landing	Quadriceps OKC knee extension.
Drills Benchmarks:	CKC exercises: lunge progression, squat progression, step-up/downs
4 months	progress with double leg / off-set foot position / single leg
Passes testing criteria -	progress single direction to multiple directions.
See next page	Balance exercises: Single leg, progress to dynamic and reactive
	Return to running program if passes benchmarks- see next page
During Landing drills :	4 months-5 months: continue with strengthening and dynamic balance
Focus on:	progress to the following exercises if clinical appropriate
1.Soft landing with knee	(see side bar)
flexion > 30 deg 2. no medial	Landing drills: Low amplitude sub-max drills Challest items landings double to aircle line items.
collapse/knee valgus	Shallow jump landings, double to single line jumps,
3. no hip IR/ pelvic drop	squat jumps progress to higher level if meets criteria (see sidebar)
4. Dynamic postural	Agility drills: Low amplitude low velocity drills:
control	skipping F/B, jogging F/B, skaters, carioca
	progress to higher level with speed and complexity (when
	appropriate)
	agility ladder drills, cutting/pivoting (changing directions),
	changing speeds, anticipated to un-anticipated



DI 11/ 0.0	
Phase IV: 6-9	Return to Higher Level Activities and Sport Phase
months Goals	 Continue to progress with strengthening, landing and agility drills to pass return to sports criteria – see testing algorithm Progress to sport specific drills Address fear avoidance beliefs by graded exercise progression, cuing, positive reinforcement, referral if necessary. Return to sports at 9-12 months if passes criteria – see testing algorithm. Sports progression may take 2-4 weeks for full clearance back to full competition
Brace	Your MD may recommend a knee sleeve or functional brace to be used until 12 months from your surgery for higher level activities
Treatment recommendations	Specific interventions and treatments will depend on the testing results. Address areas of deficits and sport specific demands.
	 Strengthening exercises (if strength scores <90%) Dynamic balance exercises if indicated (Y balance <4cm, poor control) Landing/jumping/hopping drills if limb symmetry <90% on hop test and/or faulty movement patterns (stiff knee landing, assymetrical loading, knee valgus, poor postural control. Progress agility drills Progress to sport specific exercises and drills 9 months+: possible clearance for return to sport, depending on testing -see next pages for testing algorithm
Return-to-Sports Progression: (2-4 weeks, depending on tolerance)	Step 1: 1 on 1 drills (non-contact) sport specific activities Step 2:
	1 on 1 drills (contact) full speed sport specific activities Step 3: Team scrimmage (non-contact) Step 4: Team scrimmage no restrictions Step 5: Game activities with restricted playing time Step 6: Game activities with no restrictions



ACL Return-to-Running and Return-to-Sport Testing Algorithm: Determined by: Time out from surgery, Testing performance, PT/MD approval

6 weeks

1.Knee ROM

2.Hip strength:

Abduction MMT

3.SL 30 deg Stork test

4.FOTO

8 weeks:

1.Knee ROM

2.Hip strength:

Abduction MMT/dynamometry Hip Abduction Side plank test

3.SL 30 Stork test

4.Y balance

5. Squat WB symmetry: Force plate

12 weeks (3 months)

1. Knee ROM

2.SL 60 deg Stork test

3.Hip strength:

Abduction MMT/ dynamometry / Hip Abduction Side plank test

4.Biodex test:

20 deg extension block

2 speeds: 180 deg/sec (5 reps) 300 deg/sec (30 reps)

5.Y balance test

6.Squat WB symmetry: Force plate

7.FOTO

16 weeks (4 months) - RETURN to RUNNING

Repeat previous tests not passed

For Biodex test:

20 deg extension block 3 speeds: 60 d/sec (5 reps)

180 d/sec (5 reps) 300 d/sec (30 reps)

*if adequate strength scores for return to running

(quads at least 75%, hamstrings: at least 75%)

1. Anterior lateral hop to stabilization

2.Sub-Max Jump test: no arm swing

3.Sub-Max Single Hop Assessment: no arm swing

For apprehension and control

4.Trial of running

5. Screen for fear avoidance/kinesiophobia

(ACL-RSI survey)

Return to Running Benchmarks:

1.Time: at least 4 months post-op:

2. MD / PT clearance

3. No knee joint effusion

4. ROM: limb symmetry:

extension within 5 deg flexion within 10 deg

5.Biodex: Limb symmetry of PT:

Quad: 75% Hams: 75%

6. Anterior lateral hop to stabilization drill completed with

no apprehension and good movement control

7. Assess running form: Treadmill running (sub-max at self

selected speed)
Recommendations:

1.Biodex:

Quad PT/BW: +/- 5%

Males: 95%,75%, 50% at 60,180,300deg/sec Females: 85%, 65%, 35% at 60,180,300deg/sec H/Q ratio: +/- 5%: 65%, 75%, 90% at 60,180,300deg/sec

Total work at 300 deg/sec:

Quad: limb symmetry 75% Hams: limb symmetry:75%

2. SL 60 deg stork test:

Limb symmetry: 90%

3. Hip Abductor strength: MMT 5/5 or dynamometry 90%

4. Squat WB symmetry with near equal WB

5. Y balance: Limb symmetry: < 4cm

Return to Jumping/Landing Drills Benchmarks:

1.Time: at least 4-6 months

2.MD/ PT clearance

3.No knee joint effusion

4.Biodex: Limb symmetry of PT:

Quadriceps and hamstrings: 75-85% = sub-max landing

drills

Quadriceps and hamstrings: 85-90% = max landing

drills

Minimize the following 4 variables with landing drills:

1. Stiff landing (< 30 deg knee flexion)

2. Knee valgus

3. Hip IR / pelvic drop

4. Loss of Dynamic balance



ACL Return-to-Running and Return-to-Sport Testing Algorithm

Return to sport depends on:

Timeframe from surgery Test performance MD and PT approval

24 weeks (6 months)

Repeat previous tests not passed

Biodex test: Full ROM with no ext block
 3 speed test: 60 deg/sec (5 reps),
 180 deg/sec (5 reps),

300deg/sec (30 reps

- 2. Squat WB symmetry: force plate
- 3. Landing Assessment: qualitative*
 - a. Broad jump 2D no arm swing
 - a. Land Vertical Jump 2D (front and side)
 - b. Sub-max Single leg Hop –2D (front and side) no arm swing progress to max if:

strength 90%

limited landing mechanic variable

- 4.FOTO and IKDC (Mycare)
- 5. Screen for fear avoidance/kinesiophobia

(ACL-RSI survey)

- *Landing mechanic variables at impact for potential injury risk:
 - 1. Stiff landing (< 30 deg knee flexion)
 - 2. Knee valgus
 - 3. Hip IR / pelvic drop
 - 4. Decreased dynamic balance

(poor trunk control, increased # reps to complete)

9 months- Possible return to sport

Repeat previous tests not passed

1.Biodex test: Full ROM with no ext block 3 speed test: 60 deg/sec (5 reps), 180 deg/sec (5 reps),

300deg/sec (30 reps)
2. Landing Assessment: quantitative for limb symmetry

- qualitative for landing mechanics variables
- a. Single leg hop (no arm swing) 2D (front and side)
- b. Triple hop (arm swing) 2D (front)
- c. Cross-over hop (arm swing) 2D (front)
- 3. Agility test: LEFT test components or time
- 4. FOTO and IKDC (Mycare)
- 5. screen for fear avoidance/kinesiophobia

(ACL-RSI survey)

2 year/ 5 year

IKDC (mycare)

Return to Sport Benchmarks:

1.Time: at least 9-12 months

2.MD/ PT clearance

3.No knee joint effusion

4.ROM: limb symmetry:

extension within 5 deg flexion within 10 deg

5.Biodex: Limb symmetry of PT:

Quad: 90% Hams: 90% 6.Landing Assessment:

(Single Hop/ Triple Hop/ Cross-over Hop)
Quantitative: Limb symmetry: 90%
Qualitative variables - no faulty landing
mechanics – see previous column*

7. Agility components with no compensation

8. No evidence of fear avoidance

Recommendations:

1.Biodex:

*Quad PT/BW: (+/-5%)

Males: 95%, 75%, 50% at 60, 180, 300 deg/sec Females: 85%, 65%, 35% at 60,180,300 deg/sec

H/Q ratio: (+/- 5%)

65%, 75%, 90% at 60, 180, 300 deg/sec

Hams PT/BW: (+/- 5%)

Males: 60%, 35%, 25% at 60, 180, 300 deg/sec Females: 60%, 35%, 25% at 60, 180, 300 deg/sec

<u>Total work: 300 deg/sec</u> Quads: Limb symmetry:90% Hams: Limb symmetry: 90%

2. Hip Abductor strength: MMT 5/5 or

dynamometry 90%

3.Y balance: Limb symmetry: < 4cm

4. Jump test:

Males: 90%-100% height Females: 80%-90% height

5. Single hop test:

Males: 80-90% height Females: 70-80% height



ACL Reconstruction Program References

Accelerated Rehabilitation Following ACL Reconstruction using Ipsilateral Patellar Tendon Graft Protocol printed from SportsMedRx.com, 2003

Beynnon BD, Johnson RJ. Anterior cruciate ligament injury rehabilitation in athletes. Biomechanical considerations. Sports Medicine, 1996; 22(1): 54-64

Brandsson S, Faxen E, Kartus J, et al. A prospective four to seven year follow-up after arthroscopic anterior cruciate ligament reconstruction. Scand J of Med and Science in Sports, 2001; 11(1): 23-27

Davies GJ, Heiderscheit B, Clark M. Open kinetic chain assessment and rehabilitation. Athletic Training: Sports health care perspectives, 1995; 1(4): 347-370

Davies GJ, Zillmer DA: Functional progression of exercise during rehabilitation in Knee Ligament Rehabilitation, Ellenbecker, 2000; 345-360

Delay BS, Smolinski R, Wind WM, Bowman DS. Current practices and opinions in ACL reconstruction and rehabilitation: Results of a survey of the American Orthopaedic Society for Sports Medicine (Summary). Am J of Knee Surgery, 2001; 14(2): 85-91

Fagenbaum R, Darling W. Jump landing strategies in male and female college athletes and the implications of such strategies for ACL injury. Am J of Sports Med, 2003; 31: 233-240

Falconiero RP, DiStefano VJ, Cook TM. Revascularization and ligamentization of autogenous anterior cruciate ligament grafts in humans. Arthroscopy, 1998, 14: 197-205.

Feller JA, Webster KE. A randomized comparison of patellar tendon and hamstring tendon anterior cruciate ligament reconstruction. Am J of Sports Med, 2003; 31, 564-573

Fitzgerald, KG. Open versus closed kinetic chain exercise: Issues in rehabilitation after anterior cruciate ligament reconstructive surgery. Physical Therapy, 1997; 77: 1747-1754

Heckmann TP, Noyes FR, Barber-Westin SD. Autogeneic and allogeneic anterior cruciate ligament rehabilitation. In Knee Ligament Rehabilitation, Ellenbecker 2000: 132-148

Kartus KJ, Stener S, Kohler K, et al. Is bracing after anterior cruciate ligament reconstruction necessary? A 2 year follow-up of 78 consecutive patients rehabilitated with or without a brace. Knee Surgery, Sports Traumatology, Arthroscopy, 1997; 5(3): 157-161

Kuechle DK, Pearson SE, Beach WR, et al. Allograft ACL reconstruction in patients over 40 years of age. Arthroscopy, 2002; 8: 845-853

Linn RM, Fischer DA, Smith JP, et al. Achilles tendon allograft reconstruction of the anterior cruciate ligament deficient knee. Am J of Sports Med, 1993; 21(6): 825-831

Mangine RE, Kremchek TE. Evaluation-based protocol of the anterior cruciate ligament. J of Sport Rehab, 1997; 6: 157-181



Manske RC, Prohaska D, Livermore R. Anterior Cruciate Ligament Reconstruction Using the Hamstring- Gracilis Tendon Autograph. In Postsurgical Orthopedic Sports Rehabilitation of Knee and Shoulder. Ed Manske RC. 2006: 189-206.

Martinek MV, Friederich NF. To brace or not to brace? How effective are knee braces in rehabilitation. Orthopade, 1996; 28(6): 565-570

Neitzel JA, Kernozek TW, Davies GJ: Loading response following ACL reconstruction during the parallel squat exercises. Clinical Biomechanics, 2002; 17(7): 551-554

McClure PW, Blackburn LG, Dusold C. The use of splints in the treatment of joint stiffness: biological rational and algorithm for making clinical decisions. Phy Ther,1994 74:1101-1107

Millett PJ, Wickiewicz TL, Warren RF. Motion loss after ligament injuries to the knee. Part II: Prevention and treatment. American Journal of Sports Medicine, 2001; 29: 822-828

Mikkelsen C, Werner S, Eriksson E. Closed kinetic chain alone compared to combined open and CKC exercises for quads strengthening after ACL reconstruction with respect to return to sports: a prospective matched follow-up study. Knee Sur, Sports Trau, Arthr, 2000; 8: 337-42

Muneta T, Sekiya I, Ogiuchi T, et al. Effects of aggressive early rehabilitation of the outcome of ACL reconstruction with multi-strand semitendinosus tendon. Internal Orthopaedics, 1998; 22: 352-6

Rebel M, Paessler HH. The effect of knee brace on coordination and neuronal leg muscle control: an early postoperative functional study in ACL reconstructed patients. Knee Surg, Sports Trauma, Arthro, 2001; 9(5): 272- 281

Risberg MA, Mork M, Jenssen HK, Holm I. Design and implementation of a neuromuscular training program following ACL reconstruction. J of Ortho and Sports Phy Ther, 2001; 31: 620-31

Sapega AA, Quedenfeld TC. Biophysical factors in range of motion exercises. Physician and Sports Medicine, 1981; 9: 57-65

Seo JG, Cho DY, Kim KY. Reconstruction of the anterior cruciate ligament with Achilles tendon autograft. Orthopaedic Surgery in Korea, 1993; 16(6): 719-724

Shelburne KB, Pandy MG. Determinants of cruciate-ligament loading during rehabilitation exercise. Clinical Biomechanics, 1998; 13: 403-413

Shelbourne DK, Rask BP: Controversies with anterior cruciate ligament surgery and rehabilitation. American Journal of Knee Surgery, 1998; 11(2): 136-143

Tsaklis P, Abatzides G. ACL rehabilitation program using a combined isokinetic and isotonic strengthening protocol. Isokinetics Exercise Science, 2002; 10: 211-9

Tsuda E, Okamura Y, Ishibashi Y. Techniques for reducing anterior knee symptoms after ACL using a bone-patellar tendon-bone autograft. American Journal Sports Medicine, 2001; 29: 450-456

UW Sports Medicine ACL Reconstruction in Adolescent Athlete Rehab Protocol, accessed on the website 10.24.2018

Wu GK, Ng GY, Mak AF. Effects of knee bracing on the sensorimotor function of subjects with ACL



References for early Quadriceps OKC isolated strengthening

Belloir M, Mazeas J, Traullé M, et al. Influence of the open kinetic chain on the distension of the transplant after anterior cruciate ligament surgery with hamstring graft: search for risk factors. Int J Physiother. 2020;7:256-263.

Beynnon B, Howe JG, Pope MH, Johnson RJ, Fleming BC. The measurement of anterior cruciate ligament strain in vivo. Int Orthop. 1992;16:1-12

Beynnon BD, Fleming BC, Johnson RJ, Nichols CE, Renström PA, Pope MH. Anterior cruciate ligament strain behavior during rehabilitation exercises in vivo. Am J Sports Med. 1995;23:24-34

Brinlee, A. W., Dickenson, S. B., Hunter-Giordano, A., & Snyder-Mackler, L. (2022). ACL reconstruction rehabilitation: clinical data, biologic healing, and criterion-based milestones to inform a return-to-sport guideline. *Sports Health*, *14*(5), 770-779.

Englander ZA, Garrett WE, Spritzer CE, DeFrate LE. In vivo attachment site to attachment site length and strain of the ACL and its bundles during the full gait cycle measured by MRI and high-speed biplanar radiography. J Biomech. 2020;98:109443.

Escamilla RF, Macleod TD, Wilk KE, Paulos L, Andrews JR. ACL strain and tensile forces for weight bearing and non-weight-bearing exercises after ACL reconstruction: a guide to exercise selection. J Orthop Sport Phys Ther. 2012;42:208-220

Fleming BC, Oksendahl H, Beynnon BD. Open- or closed-kinetic chain exercises after anterior cruciate ligament reconstruction? Exerc Sport Sci Rev. 2005;33:134-140

Forelli, F., Barbar, W., Kersante, G., Vandebrouck, A., Duffiet, P., Ratte, L., ... & Rambaud, A. J. (2023). Evaluation of Muscle Strength and Graft Laxity With Early Open Kinetic Chain Exercise After ACL Reconstruction: A Cohort Study. *Orthopaedic Journal of Sports Medicine*, *11*(6), 23259671231177594.

Nelson C, Rajan L, Day J, Hinton R, Bodendorfer BM. Postoperative rehabilitation of anterior cruciate ligament reconstruction: A systematic review. Sports Med Arthrosc. 2021. doi:10.1097/JSA.000000000000014

Noehren B, Snyder-Mackler L. Who's afraid of the big bad wolf? Open-chain exercises after anterior cruciate ligament reconstruction. J Orthop Sports Phys Ther. 2020. doi:10.2519/jospt.2020.0609

Perriman A, Leahy E, Semciw AI. The effect of open-versus closed-kinetic-chain exercises on anterior tibial laxity, strength, and function following anterior cruciate ligament reconstruction: A systematic review and meta-analysis. J Orthop Sports Phys Ther. 2018. doi:10.2519/jospt.2018.7656

Roldán E, Reeves ND, Cooper G, Andrews K. In vivo mechanical behaviour of the anterior cruciate ligament: a study of six daily and high impact activities. Gait Posture. 2017;58:201-207.

Taylor KA, Cutcliffe HC, Queen RM, et al. In vivo measurement of ACL length and relative strain during walking. J Biomech. 2013;46:478-483

